

WILL IT BE POSSIBLE TO MEASURE INTRINSIC GRAVITOMAGNETISM WITH LUNAR LASER RANGING?

LORENZO IORIO

INFN-Sezione di Pisa. Viale Unità di Italia 68, 70125 Bari (BA), Italy

e-mail: lorenzo.iorio@libero.it

Received 16 June 2008

Accepted 21 September 2008

Communicated by Jorge Pullin

In this note we mainly explore the possibility of measuring the action of the intrinsic gravitomagnetic field of the rotating Earth on the orbital motion of the Moon with the Lunar Laser Ranging (LLR) technique. Expected improvements in it should push the precision in measuring the Earth-Moon range to the mm level; the present-day Root-Mean-Square (RMS) accuracy in reconstructing the radial component of the lunar orbit is about 2 cm; its harmonic terms can be determined at the mm level. The current uncertainty in measuring the lunar precession rates is about 10^{-1} milliarcseconds per year. The Lense-Thirring secular, i.e. averaged over one orbital period, precessions of the node and the perigee of the Moon induced by the Earth's spin angular momentum amount to 10^{-3} milliarcseconds per year yielding transverse and normal shifts of $10^{-1} - 10^{-2}$ cm yr $^{-1}$. In the radial direction there is only a short-period, i.e. non-averaged over one orbital revolution, oscillation with an amplitude of 10^{-5} m. Major limitations come also from some systematic errors induced by orbital perturbations of classical origin like, e.g., the secular precessions induced by the Sun and the oblateness of the Moon whose modelled parts are several times larger than the Lense-Thirring signal. The present analysis holds also for the Lue-Starkman perigee precession due to the multidimensional braneworld model by Dvali, Gabadadze and Porrati (DGP); indeed, it amounts to about 5×10^{-3} milliarcseconds per year.

Keywords: Experimental studies of gravity; Moon

1. Introduction

In the framework of the linearized weak-field and slow-motion approximation of general relativity, the gravitomagnetic effects^{1,2} are induced by the off-diagonal components g_{0i} , $i = 1, 2, 3$ of the space-time metric tensor^{3,4} which are proportional to the components of the matter current density of the source $j_i = \rho v_i$.

There are essentially two types of mass currents in gravity⁵. The first type is induced by the rotation of the matter source around its center of mass and generates the intrinsic gravitomagnetic field which is closely related to the proper angular momentum $\mathbf{S}$ (i.e. spin) of the rotating body. The other type is due to the translational motion of the source and is responsible for the extrinsic gravitomagnetic field.

A debate has recently arisen concerning the possibility of measuring some extrinsic gravitomagnetic orbital effects affecting the motion of the Earth-Moon sys-

tem in the Sun's field with the Lunar Laser Ranging (LLR) technique^{6,7,8,9,10,11}. Another test of extrinsic gravitomagnetism concerning the deflection of electromagnetic waves by Jupiter in its orbital motion has been performed in a dedicated radio-interferometric experiment¹².

In this brief note we wish to consider in some details the possibility of measuring with LLR an effect induced by the intrinsic gravitomagnetic field of the spinning Earth

$$\mathbf{B}_g = \frac{G [3\mathbf{r} (\mathbf{r} \cdot \mathbf{S}) - r^2 \mathbf{S}]}{cr^5} \quad (1)$$

through the non-central, Lorentz-like acceleration

$$\mathbf{a} = -2 \left(\frac{\mathbf{v}}{c} \right) \times \mathbf{B}_g \quad (2)$$

on the orbital motion of the Moon around the Earth. In eq. (1) and eq. (2) G is the Newtonian gravitational constant, c is the speed of light in vacuum, $\mathbf{S}$ is the Earth's spin angular momentum and $\mathbf{v}$ is the velocity of the Moon with respect to the Earth. The orbital feature we are interested in consists of the Lense-Thirring precessions of the longitude of the ascending node Ω and the argument of pericentre ω of the orbit of a test particle¹³. Twenty years ago, Bertotti in Ref. 14 wrote that the Lense-Thirring effect for the Moon was, at that time, too small to be detected; according to Ciufolini¹⁰, intrinsic gravitomagnetism is still unmeasurable with the lunar orbit. Instead, the possibility of measuring it in view of the expected forthcoming improvements in LLR has recently been envisaged by Müller *et al.* in Ref. 15 and in Ref. 16; Kopeikin *et al.* in Ref. 17 retain highly plausible its measurement. An overview of other attempts to measure the Lense-Thirring effect in various Solar System scenarios with natural and artificial test particles can be found in Ref. 18. In particular, for the LAGEOS-LAGEOS II test in the gravitational field of the Earth see Ref. 19 and Ref. 20; for the perspectives on measuring the solar intrinsic gravitomagnetic field with the inner planets of the Solar System see Ref. 21; for the test performed with the Mars Global Surveyor in the field of Mars see Ref. 22 and Ref. 23. Another effect induced by the intrinsic gravitomagnetic field of the Earth is the precession of orbiting gyroscopes^{24,25} currently under measurement by the GP-B mission; see on the WEB <http://einstein.stanford.edu/>, Ref. 26 and Ref. 27.

It is interesting to note that our analysis is equally valid also for the anomalous Lue-Starkman²⁸ perigee precession predicted in the framework of the multi-dimensional braneworld model of modified gravity put forth by Dvali, Gabadadze and Porrati²⁹ (DGP) to explain the observed acceleration of the Universe without resorting to dark energy; indeed, as we will see, the magnitude of such an effect is the same as the Lense-Thirring one for the Moon. Several researchers^{28,30,31} argued that it might be possible to measure the Lue-Starkman precession with LLR in view of the expected improvements in such a technique.

The physical and geocentric orbital parameters of the Moon are listed in In Table 1.

Table 1. Physical and geocentric orbital parameters of the Moon^{32,33}. The gravity field adopted is the LP150Q solution (See on the WEB http://pds-geosciences.wustl.edu/geo/lp-l-rss-5-gravity-v1/lp_1001/sha/jgl150ql.lbl).

Parameter	Value	Units
m mass	7.349×10^{22}	kg
S proper angular momentum	2.32×10^{29}	kg m ² s ⁻¹
Gm	$4.902801076 \times 10^{12}$	m ³ s ⁻²
R radius	1.738×10^6	m
α proper angular velocity	2.66×10^{-6}	rad s ⁻¹
$\frac{C}{mR^2}$ normalized moment of inertia	0.3932	-
J_2 mass quadrupole moment	2.0326×10^{-4}	-
δJ_2 uncertainty in the mass quadrupole moment	1×10^{-8}	-
a semi-major axis	3.84400×10^8	m
I mean inclination to the Earth's equator	23.5	deg
e eccentricity	0.0549	-

2. The Lense-Thirring effect on the lunar orbit

By assuming a suitably constructed geocentric equatorial frame, it turns out that the node and the perigee of the Moon undergo the Lense-Thirring secular precessions

$$\begin{cases} \dot{\Omega}_{\text{LT}} = \frac{2GS_{\oplus}}{c^2 a^3 (1-e^2)^{3/2}} = 0.001 \text{ mas yr}^{-1}, \\ \dot{\omega}_{\text{LT}} = -\frac{6GS_{\oplus} \cos I}{c^2 a^3 (1-e^2)^{3/2}} = -0.003 \text{ mas yr}^{-1}, \end{cases} \quad (3)$$

where a, e, I are the semi-major axis, the eccentricity and the inclination to the Earth's equator of the Moon's orbit; mas yr⁻¹ stands for milliarcseconds per year. We used $S_{\oplus} = 5.85 \times 10^{33}$ kg m² s⁻¹³⁴. The Lue-Starkman²⁸ pericentre precession is just about^a ∓ 0.005 mas yr⁻¹.

Since the ratio of the mass of the Moon to that of the Earth is³⁵ $\mu = 0.0123000383$, one may argue that eq. (3), which has been derived for a test-particle like, e.g., an artificial satellite, does not apply to the Earth-Moon system. The intrinsic gravitomagnetic spin-orbit effects in the case of a two-body system with arbitrary masses m_A and m_B and spins S_A and S_B have been derived by Barker and O'Connell in Ref. 36, Damour in Ref. 37, Wex in Ref. 38; for the sake of simplicity, we will reason in terms of the node. In this case, the total node precession

^aThe minus sign is related to the standard Friedmann-Lemaître-Robertson-Walker (FLRW) branch, while the plus sign is related to the self-accelerated branch which should be able to explain the observed acceleration of the Universe without resorting to dark energy³⁰.

$\dot{\Omega}_{\text{tot}}$ accounts for the spin-orbit contributions of both bodies and also for a spin-spin term. The expression of the node precession of a body A is^{36,37,38}

$$\dot{\Omega}_A = \left(\frac{3 + x_A}{2c^2} \right) \frac{G(m_A + m_B)}{a^3(1 - e^2)^{3/2}} \frac{S_A}{m_A}, \quad x_A = \frac{m_A}{m_A + m_B}, \quad (4)$$

so that

$$\dot{\Omega}_{\text{tot}} = \dot{\Omega}_A + \dot{\Omega}_B. \quad (5)$$

Let us pose

$$m_A = m_{\oplus} \equiv M, \quad m_B = m_{\text{Moon}} \equiv m; \quad (6)$$

thus, it is possible to obtain

$$\dot{\Omega}_{\oplus} = \left(1 + \frac{3}{4}\mu \right) \frac{2GS_{\oplus}}{c^2 a^3 (1 - e^2)^{3/2}}, \quad (7)$$

$$\dot{\Omega}_{\text{Moon}} = \left(1 + \frac{3}{4\mu} \right) \frac{2GS_{\text{Moon}}}{c^2 a^3 (1 - e^2)^{3/2}}; \quad (8)$$

recall that μ is the Moon/Earth mass ratio. It results that the precession of eq. (7) is larger than the Lense-Thirring one of eq. (3) by the multiplicative factor $(1 + \frac{3}{4}\mu) = 1.0092$ yielding an error of $10^{-5} \text{ mas yr}^{-1}$, which is completely negligible (see Section 3). Concerning the precession due to the lunar spin, we have

$$\frac{\dot{\Omega}_{\text{Moon}}}{\dot{\Omega}_{\oplus}} = \left(\frac{3 + 4\mu}{4 + 3\mu} \right) \frac{1}{\mu} \frac{S_{\text{Moon}}}{S_{\oplus}} = 2 \times 10^{-3}, \quad (9)$$

i.e. it is of the order of $2 \times 10^{-6} \text{ mas yr}^{-1}$, which is negligible as well. The amplitude of the spin-spin term is proportional to^{36,37,38}

$$\dot{\Omega}_{\text{SS}} \propto -\frac{3}{2c^2} \sqrt{\frac{GM(1 + \mu)}{a^7}} \frac{1}{(1 - e^2)^2} \frac{S_{\oplus}}{M} \frac{S_{\text{Moon}}}{m} = 6 \times 10^{-9} \text{ mas yr}^{-1}. \quad (10)$$

Thus, we can conclude that the Lense-Thirring approximation is fully adequate for the Earth-Moon system.

3. Some sources of error

Let us now examine some sources of systematic errors. In regard to the potentially corrupting action of the mismodelling in the even ($\ell = 2, 4, 6, \dots$) zonal ($m = 0$) harmonic coefficients J_{ℓ} of the multipolar expansion of the Newtonian part of the Earth's gravitational potential, which is not the most important source of aliasing precessions in the case of the Moon³², only δJ_2 would be of some concern. Indeed, the mismodelled secular precessions induced by it on the lunar node and perigee amount to^b $-2.67 \times 10^{-4} \text{ mas yr}^{-1}$ and $5.3 \times 10^{-4} \text{ mas yr}^{-1}$, respectively; the impact of the other higher degree even zonals is negligible being $\leq 10^{-8} \text{ mas yr}^{-1}$.

^bThe calibrated errors δJ_{ℓ} of the EIGEN-CG01C Earth gravity field solution³⁹ were used.

As in the case of the spins, also the asphericity of the Moon has to be taken into account^{36,38} according to

$$\dot{\Omega}_{J_2^{\text{Moon}}} = -\frac{3}{2} \frac{n_{\text{Moon}} \cos F J_2^{\text{Moon}}}{(1-e^2)^2} \left(\frac{R_{\text{Moon}}}{a} \right)^2, \quad (11)$$

where $n_{\text{Moon}} = \sqrt{GM(1+\mu)/a^3}$ is the lunar mean motion and F is the angle between the orbital angular momentum and the Moon's spin angular momentum $\mathbf{S}_{\text{Moon}}$; it is about 3.61 deg since the spin axis of the Moon is tilted by 1.54 deg to the ecliptic and the orbital plane has an inclination of 5.15 deg to the ecliptic³³. Table 1 and eq. (11) yield a mismodelled node precession due to δJ_2^{Moon} of about $0.006 \text{ mas yr}^{-1}$, which is 6 times larger than the Lense-Thirring rate. For other sources of systematic errors induced by gravitational and even non-gravitational⁴⁰ perturbations see Ref. 32 and references therein, especially Ref. 41. Among the N-body gravitational perturbations, the largest ones are due to the Sun's attraction. In order to get an order-of-magnitude evaluation of their mismodelling, let us note that some of such effects are proportional to $n_{\oplus}^2/n_{\text{Moon}}$; e.g. the node rate, referred to the equator, is⁴²

$$\dot{\Omega}^{\odot} = \frac{3G\mathfrak{M}_{\odot} \cos I}{4a_{\oplus}^3 n_{\text{Moon}}} \left(\frac{3}{2} \sin^2 \varepsilon - 1 \right) \approx -5 \times 10^7 \text{ mas yr}^{-1}, \quad (12)$$

where $\varepsilon = 23.439 \text{ deg}$ is the obliquity of the ecliptic. Since³⁵ $\delta G\mathfrak{M}_{\odot} = 5 \times 10^{10} \text{ m}^3 \text{ s}^{-2}$ and⁴³ $\delta GM = 8 \times 10^5 \text{ m}^3 \text{ s}^{-2}$, we can assume a bias of $\approx 0.07 \text{ mas yr}^{-1}$ which is 70 times larger than the Lense-Thirring precession.

Let us, now, consider the precision of LLR in reconstructing the lunar orbit with respect to the Lense-Thirring effect. Concerning the precision in measuring the lunar precession rates, it amounts to about^{44,45,15,16} 0.1 mas yr^{-1} , i.e. it is two orders of magnitude larger than the Lense-Thirring precessions of eq. (3). The orbital perturbations experienced by a test particle are usually decomposed along three orthogonal directions of a frame co-moving with it; they are named radial R (along the radius vector), transverse T (orthogonal to the radius vector, in the osculating orbital plane) and normal N (along the orbital angular momentum, out of the osculating orbital plane). According to, e.g., Ref. 46, the R - T - N perturbations can be expressed in terms of the shifts in the Keplerian orbital elements as

$$\begin{cases} \Delta R = \sqrt{(\Delta a)^2 + \frac{[(e\Delta a + a\Delta e)^2 + (ae\Delta \mathcal{M})^2]}{2}}, \\ \Delta T = a\sqrt{1 + \frac{e^2}{2}} \left[\Delta \mathcal{M} + \Delta \omega + \cos I \Delta \Omega + \sqrt{(\Delta e)^2 + (e\Delta \mathcal{M})^2} \right], \\ \Delta N = a\sqrt{1 + \frac{e^2}{2}} \left[\frac{(\Delta I)^2}{2} + (\sin I \Delta \Omega)^2 \right], \end{cases} \quad (13)$$

where $\mathcal{M}$ is the mean anomaly. The lunar Lense-Thirring shifts after one year are, thus

$$\begin{cases} \Delta R_{LT} = 0, \\ \Delta T_{LT} = a\sqrt{1 + \frac{e^2}{2}} (\Delta\omega_{LT} + \cos I \Delta\Omega_{LT}) = -0.38 \text{ cm}, \\ \Delta N_{LT} = a\sqrt{1 + \frac{e^2}{2}} \sin I \Delta\Omega_{LT} = 0.07 \text{ cm}. \end{cases} \quad (14)$$

It is important to note that there is no Lense-Thirring secular signature in the Earth-Moon radial motion on which all of the efforts of LLR community have been concentrated so far. It can be shown that a short-period, i.e. not averaged over one orbital revolution, radial signal exists; it is proportional to

$$\Delta r \propto \frac{2GS_{\oplus}}{c^2 na^2} = 2 \times 10^{-5} \text{ m}, \quad (15)$$

which is too small to be detected since the present-day accuracy in estimating the amplitudes of radial harmonic signals is of the order of mm⁶. Major limitations come from the post-fit Root-Mean-Square (RMS) accuracy with which the lunar orbit can be reconstructed; the present-day accuracy is about 2 cm in the radial direction R along the centers-of-mass of the Earth and the Moon¹⁶. Improvements in the precision of the Earth-Moon ranging of the order of 1 mm are expected in the near future with the APOLLO program^{47,48}. Recently, sub-centimeter precision in determining range distances between a laser on the Earth's surface and a retro-reflector on the Moon has been achieved⁴⁹. However, it must be considered that the RMS accuracy in the T and N directions is likely worse than in R .

4. Conclusions

In this note we have examined the possibility of measuring the action of the intrinsic gravitomagnetic field of the spinning Earth on the lunar orbital motion with the LLR technique. After showing that the Lense-Thirring approximation is adequate for the Earth-Moon system, we found that the Lense-Thirring secular precessions of the Moon's node and the perigee induced by the Earth's spin angular momentum are of the order of 10^{-3} mas yr⁻¹ corresponding to transverse and normal secular shifts of $10^{-1} - 10^{-2}$ cm yr⁻¹. The intrinsic gravitomagnetic field of the Earth does not secularly affect the radial component of the Moon's orbit; a short-period, i.e. not averaged over one orbital revolution, radial oscillation is present, but its amplitude is of the order of 10^{-5} m. The current RMS accuracy in reconstructing the lunar orbit is of the order of cm in the radial direction; the harmonic components can be determined at the mm level. Forthcoming expected improvements in LLR should allow to reach the mm precision in the Earth-Moon ranging. The present-day accuracy in measuring the lunar precessional rate is of the order of 10^{-1} mas yr⁻¹. Major limitations come also from some orbital perturbations of classical origin like,

e.g., the secular node precessions induced by the Sun and the oblateness of the Moon which act as systematic errors and whose mismodelled parts are up to 70 times larger than the Lense-Thirring effects. As a consequence of our analysis, we are skeptical concerning the possibility of measuring intrinsic gravitomagnetism with LLR in a foreseeable future. The same conclusion holds also for the Lue-Starkman perigee precession predicted in the framework of the multidimensional braneworld DGP model of modified gravity; indeed, it is as large as the Lense-Thirring one for the Moon.

Acknowledgments

I thank T. Murphy and J.G. Williams (NASA-JPL) for useful information and material supplied.

References

1. M.L. Ruggiero and A. Tartaglia, *N. Cim. B* **117** (2002) 743.
2. G. Schäfer, *Gen. Relativ. Gravit.* **36** (2004) 2223.
3. B. Mashhoon, Gravitoelectromagnetism in *Reference Frames and Gravitomagnetism* eds. J.-F. Pascual-Sánchez, L. Floría, A. San Miguel and F. Vicente (World Scientific, 2001), p. 121.
4. B. Mashhoon, Gravitoelectromagnetism: A Brief Review in *The Measurement of Gravitomagnetism: A Challenging Enterprise* ed. L. Iorio (NOVA publishers, 2007), p. 29.
5. S.M. Kopeikin, *Int. J. Mod. Phys. D* **15** (2006) 305.
6. T.W. Murphy, K. Nordtvedt and S.G. Turyshev, *Phys. Rev. Lett.* **98** (2007a) 071102.
7. S.M. Kopeikin, *Phys. Rev. Lett.* **98** (2007) 229001.
8. T.W. Murphy, K. Nordtvedt and S.G. Turyshev, *Phys. Rev. Lett.* **98** (2007b) 229002.
9. M. Soffel, S. Klioner, J. Müller and L. Biskupek, *Phys. Rev. D* **78** (2008) 024033.
10. I. Ciufolini, arXiv:0809.3219v1 [gr-qc] (2008).
11. S.M. Kopeikin, arXiv:0809.3392v1 [gr-qc] (2008).
12. E.B. Fomalont and S.M. Kopeikin, Radio interferometric tests of general relativity in *A Giant Step: from Milli- to Micro-arcsecond Astrometry Proceedings IAU Symposium No. 248, 2007* eds. W.J. Jin, I. Platais and M.A.C. Perryman (Cambridge University Press, 2008), p. 383.
13. J. Lense J and H. Thirring, *Phys. Z.* **19** (1918) 156
14. B. Bertotti, *Celest. Mech.* **44** (1988) 31.
15. J. Müller, J.G. Williams, S.G. Turyshev and P.J. Shelus, Potential Capabilities of Lunar Laser Ranging for Geodesy and Relativity in *Dynamic Planet* eds. P. Tregoning and C. Rizos (Springer Verlag, 2007), p. 903.
16. J. Müller, J G Williams and S.G. Turyshev, Lunar Laser Ranging Contributions to Relativity and Geodesy in *Lasers, Clocks and Drag-Free Control* eds. H. Dittus, C. Lämmerzahl and S.G. Turyshev (Springer Verlag, 2008), p. 457.
17. S. Kopeikin, E. Pavlis, D. Pavlis, V.A. Brumberg, A. Escapa, J. Getino, A. Gusev, J. Müller, W.-T. Ni and N. Petrova, *Adv. Sp. Res.* **42** (2008) 1378.
18. L. Iorio, (ed.) *The Measurement of Gravitomagnetism: A Challenging Enterprise* (NOVA publishers, 2007).
19. I. Ciufolini and E.C. Pavlis, *Nature* **431** (2004) 958.
20. J.C. Ries, R.J. Eanes and M.M. Watkins, Confirming the Frame-Dragging Effect with

- Satellite Laser Ranging, Abstract presented at 16th International Laser Ranging Workshop, (Poznań, 13-17 October, 2008).
21. L. Iorio, *Scholarly Research Exchange* **2008** (2008) 105235.
 22. L. Iorio, *Class. Quantum Grav.* **23** (2006) 5451.
 23. K. Krogh, *Class. Quantum Grav.* **24** (2007) 5709.
 24. G.E. Pugh, *WSEG Research Memorandum No. 11* (1959) Reprinted in *Nonlinear Gravitodynamics. The Lense–Thirring Effect* eds. R.J. Ruffini and C. Sigismondi (World Scientific, 2003), p. 414.
 25. L. Schiff, *Am. J. of Phys.* **28** (1960) 340.
 26. C.W.F. Everitt, The Gyroscope Experiment I. General Description and Analysis of Gyroscope Performance, in *Proc. Int. School Phys. “Enrico Fermi” Course LVI* ed. B. Bertotti (New Academic Press, 1974), p. 331.
 27. C.W.F. Everitt *et al.* Gravity Probe B: Countdown to Launch, in *Gyros, Clocks, Interferometers...: Testing Relativistic Gravity in Space* eds. C Lämmerzahl, C W F Everitt and F W Hehl (Springer Verlag, 2001), p. 52.
 28. A. Lue and G. Starkman, *Phys. Rev. D* **67** 2003 064002.
 29. G. Dvali, G. Gabadadze and M. Porrati, *Phys. Lett. B* **485** 2000 208.
 30. G. Dvali, A. Gruzinov and M. Zaldarriaga, *Phys. Rev. D* **68** 2003 024012.
 31. S.G. Turyshev, *Annu. Rev. Nucl. Part. Sci.* **58** 2008 207.
 32. J.G. Williams and J.O. Dickey, Lunar Geophysics, Geodesy, and Dynamics, in *Proc. of the 13th International Workshop on Laser Ranging “Toward Millimeter Accuracy”* eds. R. Noomen, S. Klosko, C. Noll and M. Pearlman (NASA/CP-2003-212248, 2003) http://cdsis.nasa.gov/lw13/docs/papers/sci_williams_1m.pdf
 33. R.B. Roncoli, *Lunar Constants and Models Document* (JPL Technical Document D-32296, 2005).
 34. D.D. McCarthy and G. Petit, *IERS Conventions (2003)* (Frankfurt am Main: Verlag des Bundesamtes für Kartographie und Geodäsie, 2004), p. 106.
 35. E.M. Standish, (JPL IOM 312-F,1998).
 36. B.M. Barker and R.F. O’Connell, *Phys. Rev. D* **12** (1975) 329.
 37. T. Damour and G. Schäfer, *Nuovo Cimento B* **101** (1988) 127.
 38. N. Wex, *Class. Quantum Grav.* **12** (1995) 983.
 39. Ch. Reigber *et al.* Scientific Technical Report STR06/07, GeoForschungsZentrum Potsdam (2006)
 40. D. Vokrouhlický, *Icarus* **126** (1999) 293.
 41. M. Chapront-Touzé and J. Chapront, *Astron. Astrophys.* **124** (1983) 50.
 42. B.D. Tapley, B.E. Schutz and G.H. Born, *Statistical Orbit Determination* (Elsevier Academic Press, 2004).
 43. E. Groten, Report of the IAG. Special Commission SC3, Fundamental Constants, *XXII IAG General Assembly* (1999).
 44. J. Müller, M. Schneider, M. Soffel and H. Ruder, *Astrophys. J.* **382** (1991) L101.
 45. J.G. Williams, Newhall X X and J.O. Dickey, *Phys. Rev. D* **53** (1996) 6730.
 46. D.C. Christodoulidis, D.E. Smith, R.G. Williamson and S.M. Klosko, *J. Geophys. Res.* **93**(B6) (1988) 6216.
 47. J.G. Williams, S.G. Turyshev and T.W. Murphy, *Int. J. Mod. Phys. D* **13** (2004) 567.
 48. T.W. Murphy *et al.* *Pub. Astron. Soc. Pacif.* **120** (2008) 20.
 49. J. Battat *et al.* Abstract #K12.003 presented at *American Physical Society, APS April Meeting* (14-17 April, 2007).